arXiv:1704.01908v2 [math.NT] 24 Apr 2017

PRIMES IN HIGHER-ORDER PROGRESSIONS ON AVERAGE

NIANHONG ZHOU

Abstract. In this paper, we establish some theorems on the distribution of primes in higher-order progressions on average.

1. Introduction

The Bateman-Horn conjecture [1] suggests that if $x^\ell + u \in \mathbb{Z}[x]$ be irreducible polynomial with u be an even number and the degree $\ell \geq 1$. Then

$$\sum_{m\leq X} \Lambda(m)\Lambda(m^\ell + u) \sim \prod_p \left\{ \left(1-\frac{1}{p}\right)^{-2} \left(1-\frac{n_\ell(p,u)}{p}\right)\right\} X, \tag{1.1}$$

where Λ denotes the von Mangoldt function, the product being taken over all primes and $n_\ell(p,u)$ being the number of solutions of the congruence $x(x^\ell+u) \equiv 0 \pmod p$.

If $\ell = 1$, the asymptotic formula in (1.1) is the twin prime conjecture. However, even the simple case seems beyond the current approach. In 1970, Lavrik [2] proved that if $\ell = 1$, then given any $A > 0$, (1.1) holds for all even integer $u \geq 1$ not exceeding X with at most $O\left(X(\log X)^{-A}\right)$ exceptions.

In [3], S. Baier and L. Zhao established certain theorems for the Bateman-Horn conjecture for quadratic polynomials on average. Their main result states the following. Given $A, B > 0$, we have, for $x^2(\log x)^{-A} \leq y \leq x^2$,

$$\sum_{k\leq y, \mu^2(k)=1} \left| \sum_{n\leq x} \Lambda(n^2+k) - \mathfrak{S}(k)x \right|^2 \ll yx^2(\log x)^{-B},$$

where

$$\mathfrak{S}(k) = \prod_{p>2} \left(1 - \frac{1}{p-1}\left(\frac{-k}{p}\right)\right)$$

with $\left(\frac{-k}{p}\right)$ being the Legendre symbol. In [4], F. Too and L. Zhao established similar results for the cubic cases.

In this paper, we shall study the asymptotic formula in (1.1) on average. Our main results are as follows.

Theorem 1.1. *Let integer $\ell \geq 2$. For any $A > 0$, there exists a $B = B_\ell(A)$ such that*

$$\sum_{1\leq u\leq y} \left| \sum_{m\leq \sqrt[\ell]{X}} \Lambda(m^\ell+u)\Lambda(m) - \mathfrak{S}_\ell(u)\sqrt[\ell]{X} \right|^2 \ll \frac{yX^{\frac{2}{\ell}}}{\log^A X}$$

2010 *Mathematics Subject Classification.* Primary: 11P55, Secondary: 11N13, 11N32.

Key words and phrases. primes represented by polynomials; Bateman-Horn conjecture; circle method.

holds for any $y \in (X^{1-\frac{1}{2\ell}}(\log X)^B, X]$, *where*

$$\mathfrak{S}_\ell(u) = \prod_{p|u} \frac{p - \varrho_\ell(p,u)}{p - 1 - \varrho_\ell(p,u)} \prod_p \left(1 - \frac{\varrho_\ell(p,u) - 1}{p-1} - \frac{\varrho_\ell(p,u)}{(p-1)^2}\right),$$

the product being taken over all primes and $\varrho_\ell(p,u)$ *being the number of solutions of the congruence* $x^\ell + u \equiv 0 \pmod p$.

By similar arguments, we have the following theorem which improves the results in [3] and [4].

Theorem 1.2. *Let integer* $\ell \geq 2$. *For any* $A > 0$ *and* $\varrho_\ell(p,u)$ *as defined in the Theorem 1.1. Then there exists a* $B' = B'_\ell(A)$ *such that*

$$\sum_{1 \leq u \leq y} \left| \sum_{m \leq \sqrt[\ell]{X}} \Lambda(m^\ell + u) - \mathfrak{S}'_\ell(u) \sqrt[\ell]{X} \right|^2 \ll \frac{y X^{\frac{2}{\ell}}}{\log^A X}$$

holds for any $y \in (X^{1-\frac{1}{\ell}}(\log X)^{B'}, X]$ *with*

$$\mathfrak{S}'_\ell(u) = \prod_p \left(1 - \frac{\varrho_\ell(p,u) - 1}{p-1}\right),$$

the product being taken over all primes.

The primary technique used in the proof of Theorem 1.1 is the circle method and the use of a variant of Weyl's inequality. The main difficulty in this application of the circle method is with the singular series. As for the asymptotic conjecture (1.1), the coefficient $\mathfrak{S}_\ell(u)$ involves the use of Dedekind zeta functions associated to suitable algebraic number fields of the form $\mathbb{Q}[\sqrt[\ell]{u}]$. On the other hand, let p, q denotes prime and we observe that left of (1.1) means that one can give an estimate for

$$\#\{q \in \mathbb{N} : q^\ell + u = p, q \leq X\} = \#\{q \in \mathbb{N} : p - q^\ell = u, q \leq X\}.$$

Which similar with the Hardy-Littlewood conjecture [5], say every sufficiently large number is either a ℓ-*th* power or a sum of a prime number and a ℓ-*th* power, for $\ell = 2, 3$. When the circle method used, in fact there is no big difference between them. Therefore when $\ell \geq 2$, the singular series similar as the singular series of Zaccagnini [6], which first give a crude estimates for the kinds of singular series. In [7], Kawada announced that he could get an asymptotic formula for the number of representations of numbers as the sum of a prime and a ℓ-*th* power on average, and give a detailed proof in [8] by use the analytic properties of the Dedekind zeta function. Based on this results and under Generalized Riemann Hypothesis, Brödern [9] give an asymptotic formula for the number of representations of numbers as the sum of a prime and a ℓ-*th* power of a prime on average.

Furthermore, combine with the work of Perelli, Zaccagnini [10] and Bauer [11], we can have a good treatment for the minor arcs. Hence get the proof of our main theorem.

Notation. Notation is standard or otherwise introduced when appropriate. The symbols $\mathbb{Z}$ and $\mathbb{Q}$ denote the integers and the rational numbers, respectively. $e(z) = e^{2\pi i z}$, the letter p always denotes a prime. The symbol $\mathbb{Z}_q$ represents shorthand for the groups $\mathbb{Z}/q\mathbb{Z}$. Also, the shorthand for the multiplicative group reduced residue classes $(\mathbb{Z}/q\mathbb{Z})^*$ is $\mathbb{Z}_q^*$. Denote by φ and Λ the Euler and von Mangoldt functions, respectively. For a large number X, denote $L = \log X$. For the sake of simplicity, we set

$$I_\ell(\alpha, z) = \sum_{z < m^\ell \leq 2^\ell z} e\left(m^\ell \alpha\right), \quad J_\ell(\alpha, z) = \sum_{z < m^\ell \leq 2^\ell z} \Lambda(m) e\left(m^\ell \alpha\right),$$

$$I(\alpha,z)=\sum_{m\le 2^{\ell}z}e\left(-m\alpha\right)\text{ and }J(\alpha,z)=\sum_{m\le 2^{\ell}z}\Lambda(m)e\left(-m\alpha\right).$$

Otherwise, we set

$$\lambda(q,u)=\frac{1}{q}\sum_{a\in\mathbb{Z}_q^*}\sum_{h\in\mathbb{Z}_q}e\left(\frac{a(h^{\ell}+u)}{q}\right)\text{ and }A(q,u)=\frac{1}{\varphi(q)}\sum_{a\in\mathbb{Z}_q^*}\sum_{h\in\mathbb{Z}_q^*}e\left(\frac{a(h^{\ell}+u)}{q}\right).$$

It is easily seen that both $\lambda(q,u)$ and $A(q,u)$ are multiplicative function in terms of positive integer q. It is obvious that

$$\lambda(q,u)=\prod_{p|q}\left(\varrho_{\ell}(p,u)-1\right)$$

and

$$A(q,u)=\frac{q}{\varphi(q)}\prod_{p|q(q,u)^{-1}}\left(\varrho_{\ell}(p,u)-1+\frac{1}{p}\right)\prod_{p|(u,q)}\left(\varrho_{\ell}(p,u)-2+\frac{1}{p}\right)$$

when q is square-free. Also, for any $z\ge 1$, we always set

$$\mathfrak{S}'_{\ell}(u,z)=\sum_{q\le z}\frac{\mu(q)}{\varphi(q)}\lambda(q,u),\ \mathfrak{S}_{\ell}(u,z)=\sum_{q\le z}\frac{\mu(q)}{\varphi(q)}A(q,u)$$

$$\mathfrak{P}_{\ell}(u,z)=\prod_{p|u,p\le z}\frac{p-\varrho_{\ell}(p,u)}{p-1-\varrho_{\ell}(p,u)}\prod_{p\le z}\left(1-\frac{\varrho_{\ell}(p,u)-1}{p-1}-\frac{\varrho_{\ell}(p,u)}{(p-1)^2}\right)$$

and

$$\mathfrak{P}'_{\ell}(u,z)=\prod_{p\le z}\left(1-\frac{\varrho_{\ell}(p,u)-1}{p-1}\right).$$

2. Preliminary lemma

We shall need the following well-known results in analytic number theory which need in the sequel.

Lemma 2.1. *Let $1\le z\le y$, $v\in\mathbb{Z}\setminus\{0\}$ and integer $\ell\ge 2$. Then we have*

$$\sum_{y<n\le 2y}\left|\mathfrak{S}'_{\ell}(nv,z)-\mathfrak{S}'_{\ell}(nv)\right|^2\ll_{v,\ell}yz^{-1/(2000\ell^2)}.$$

Proof. This is due from Theorem 1, Corollary 1 and Corollary 2 of [8]. □

Lemma 2.2. *Let $|u|\ge 1$, $\ell\in\mathbb{Z}_{\ge 1}$ and $x^{\ell}+u$ is irreducible over $\mathbb{Q}[x]$. Then we have*

$$\left|\sum_{p}\frac{\varrho_{\ell}(p,u)-1}{p}\right|\le O_{\ell}\left(1\right)+4\ell\log\log(2\left|u\right|).$$

Proof. Let D_u be the discriminant of $\mathbb{Q}[\sqrt[\ell]{u}]$. It is easily seen that $|D_u|\le\ell^{\ell}\left|u\right|^{\ell-1}$. Hence by Landau prime ideal theorem (see [12, Theorem 5.33]) and part integration we have

$$\begin{aligned}\left|\sum_{p}\frac{\varrho_{\ell}(p,u)-1}{p}\right|&\le(\ell-1)\sum_{p\le y}\frac{1}{p}+\left|\sum_{p>y}\frac{\varrho_{\ell}(p,u)-1}{p}\right|\\&\le(\ell-1)\log\log y+O\left(\sqrt{D_u}\exp\left(-c_{\ell}\sqrt{\log y}\right)\log y\right),\end{aligned}$$

where c_ℓ is an absolute constant depend only on ℓ. Setting $y = \exp((\log(2|u|))^4)$ we obtain that

$$\left|\sum_p \frac{\varrho_\ell(p,u)-1}{p}\right| \le 4\ell \log\log(2|u|) + O_\ell\left(1 + D_u^{\frac{1}{2}}|2u|^{-c_\ell \log|2u|}(\log|2u|)^4\right).$$

Thus if $|u| \ge \exp(\ell/c_\ell)$, then we obtain that

$$\left|\sum_p \frac{\varrho_\ell(p,u)-1}{p}\right| \le 4\ell \log\log(2|u|) + O_\ell(1).$$

Thus we get the proof of the lemma. □

Lemma 2.3. *Let $\alpha = a/q + \beta$, $|\beta| \le X^{-1}L^B$ and $q \le L^B$, then*

$$R_\ell(\alpha, z) := J_\ell(\alpha, z) - I_\ell(\beta, z)B_\ell(q,a)/\varphi(q) \ll z^{1/\ell}L^{-10B},$$
$$R'_\ell(\alpha, z) := I_\ell(\alpha, z) - I_\ell(\beta, z)B'_\ell(q,a)/q \ll z^{1/\ell}L^{-10B},$$

where

$$B_\ell(q,a) = \sum_{h\in\mathbb{Z}_q^*} e\left(ah^\ell/q\right) \text{ and } B'_\ell(q,a) = \sum_{h\in\mathbb{Z}_q} e\left(ah^\ell/q\right).$$

Proof. It is easily seen that

$$J_k(\alpha, z) = \sum_{h\in\mathbb{Z}_q^*} e\left(\frac{ah^{1/\ell}}{q}\right) \sum_{\substack{z^{1/\ell}<m\le 2z^{1/\ell}\\ m\equiv h \bmod q}} \Lambda(m)e(m^k\beta) + O(\log q \log z)$$
$$= \frac{1}{\varphi(q)} \sum_{h\in\mathbb{Z}_q^*} e\left(\frac{ah^\ell}{q}\right) \sum_{z^{1/\ell}<m\le 2z^{1/\ell}} e\left(m^\ell\beta\right) + R_\ell(\alpha, z),$$

by part summation and where

$$R_\ell(\alpha, z) = \varphi(q)(1+|\beta|z) \max_{h\in\mathbb{Z}_q^*} \max_{z^{1/\ell}<x\le 2z^{1/\ell}} \left|\sum_{\substack{m\le x\\ m\equiv h \bmod q}} \Lambda(m) - x/\varphi(q)\right|$$

Then by [12, Corollary 5.29], we get the estimate of $R_\ell(\alpha, z)$. The estimate of $R'_\ell(\alpha, z)$ is similar and we omit its detail. □

Lemma 2.4. *Let $\alpha = a/q + \lambda$ with $(a,q) = 1$ and $|\lambda| \le q^{-2}$. Then for each integer $\ell \ge 2$ and any $A > 0$ there exists a $B_{\mathfrak{m},\ell}(A) > 0$ such that for $B \ge B_{\mathfrak{m},\ell}(A)$ the estimate*

$$\int_y^{2y} \mathrm{d}t \left|\sum_{t<m^\ell\le t+H} \Lambda(m)e(m^\ell a/q)\right|^2 \ll_{\ell,A} H^2 y^{\frac{2}{\ell}-1}L^{-A-2}$$

holds for $L^B < q \le HL^{-B}, y^{1-1/(2\ell)}L^B < H \le y$.

Proof. This is quoted from [11, Lemma 3.3]. □

Lemma 2.5. *Let $q \ge L^A$ with $(a,q) = 1$. Then for each integer $\ell \ge 2$, there exists a $B'_{\mathfrak{m},\ell}(A) > 0$ such that for $B \ge B'_{\mathfrak{m},\ell}(A)$ the estimate*

$$W_\ell(y, H) := \int_y^{2y} \mathrm{d}t \left|\sum_{t<m^\ell\le t+H} e(m^\ell a/q)\right|^2 \ll_{\ell,A} H^2 y^{\frac{2}{\ell}-1}L^{-A-2}$$

holds for $L^B < q \le HL^{-B}, y^{1-1/(2\ell)}L^B < H \le y$.

Proof. It is easily seen that

$$\begin{aligned}
W_\ell(y,H) &= \sum_{\substack{y<m_1^\ell,m_2^\ell\le 2y+H\\ |m_1^\ell-m_2^\ell|\le H}}\sum e\left((m_1^\ell-m_2^\ell)a/q\right)\int_{\max(m_1^\ell-H,m_2^\ell-H)}^{\min(m_1^\ell,m_2^\ell)} \mathrm{d}x\\
&= \sum_{\substack{y<m_1^\ell,m_2^\ell\le 2y+H\\ |m_1^\ell-m_2^\ell|\le H}}\sum e\left((m_1^\ell-m_2^\ell)a/q\right)\left(H-|m_1^\ell-m_2^\ell|\right)\\
&= \sum_k\sum_n \left(H-|P_\ell(n,k)|\right)\mathbf{1}_{\substack{y<(n+k)^\ell,n^\ell\le 2y+H\\ |(n+k)^\ell-n^\ell|\le H}} e\left(P_\ell(n,k)a/q\right)\\
&\ll \sum_{k\le Hy^{1/\ell-1}} H \max_{y^{1/\ell}<x\le 2y^{1/\ell}}\left|\sum_{y^{1/\ell}\le n\le x} e\left(P_\ell(n,k)a/q\right)\right|\\
&\ll \sum_{k\le Hy^{1/\ell-1}} H \max_{y^{1/\ell}/2<x\le 2y^{1/\ell}}\left|\sum_{n\le x} e\left(P_\ell(n,k)a/q\right)\right|,
\end{aligned}$$

where $P_\ell(n,k)=(n+k)^\ell-n^\ell$, then by [10, Lemma], we have

$$W_\ell(y,H)\ll_{\ell,A} H^2y^{\frac{2}{\ell}-1}\left(\left(\frac{L^A}{q}\right)^{2^{-1+\ell}}+\left(\frac{q}{Hy^{1/\ell}}\right)^{2^{-1+\ell}}+\left(\frac{1}{L^A}\right)^{2^{-1+\ell}}\right)$$

holds for any $A>(\ell+1)^2$. By setting $B'_{\mathfrak{m},\ell}(A)=2^\ell\ell^4A$, we obtain the proof of the lemma. □

3. The proof of the main results

We first denote

$$S_\ell(y,X)=\sum_{1\le u\le y}\left|\sum_{m^\ell\le X}\Lambda(m^\ell+u)\Lambda(m)-\mathfrak{S}_\ell(u)\sum_{m^\ell\le X}1\right|^2$$

and

$$S'_\ell(y,X)=\sum_{1\le u\le y}\left|\sum_{m^\ell\le X}\Lambda(m^\ell+u)-\mathfrak{S}'_\ell(u)\sum_{m^\ell\le X}1\right|^2.$$

Then, by sum over dyadic intervals process one has

$$\begin{aligned}
S_\ell(y,X) &\ll \sum_{1\le u\le y}\left|\sum_{m^\ell\le XL^{-B}}\Lambda(m^\ell+u)\Lambda(m)-\mathfrak{S}_\ell(u)\sum_{m^\ell\le XL^{-B}}1\right|^2\\
&\quad+\sum_{1\le u\le y}\left|\sum_{XL^{-B}<m^\ell\le X}\Lambda(m^\ell+u)\Lambda(m)-\mathfrak{S}_\ell(u)\sum_{XL^{-B}<m^\ell\le X}1\right|^2\\
&\ll yX^{2/\ell}L^{-B}+BL\sup_{X/L^B\le z\le X/2^\ell}F_\ell(y,z), \qquad (3.1)
\end{aligned}$$

where $B \geq 2$ and

$$F_\ell(y,z) = \sum_{u\leq y}\left|\sum_{z<m^\ell\leq 2^\ell z}\big(\Lambda(m^\ell+u)\Lambda(m)-\mathfrak{S}_\ell(u)\big)\right|^2.$$

Similarly, we have

$$S'_\ell(y,X) \ll yX^{2/\ell}L^{-B} + BL\sup_{X/L^B\leq z\leq X/2^\ell} F'_\ell(y,z) \tag{3.2}$$

for any $B \geq 2$, where

$$F'_\ell(y,z) = \sum_{u\leq y}\left|\sum_{z<m^\ell\leq 2^\ell z}\big(\Lambda(m^\ell+u)-\mathfrak{S}'_\ell(u)\big)\right|^2.$$

We define the major arcs as

$$J_{q,a} = \Big(a/q - L^{B^2}/X, a/q + L^{B^2}/X\Big], \tag{3.3}$$

where $1 \leq a \leq q$. It is obvious that the interval $J_{q,a}$ are pairwise disjoint. Setting

$$\mathfrak{M} = \bigcup_{q\leq L^{B^2}}{\bigcup_{1\leq a\leq q}}^{*} J_{q,a} \quad \text{and} \quad \mathfrak{m} = \Big(L^{B^2}/X, 1+L^{B^2}/X\Big] \setminus \mathfrak{M}, \tag{3.4}$$

where $*$ means that $(a,q)=1$. Application of the circle method gives

$$\sum_{z<m^\ell\leq 2^\ell z}\Lambda(m^\ell+u)\Lambda(m) = \left\{\int_{\mathfrak{M}}+\int_{\mathfrak{m}}\right\} J(\alpha,z)J_\ell(\alpha,z)e(u\alpha)\,\mathrm{d}\alpha.$$

Therefore,

$$\begin{aligned} F_\ell(y,z) &= \sum_{u\leq y}\left|\sum_{z<m^\ell\leq 2^\ell z}\big(\Lambda(m^\ell+u)\Lambda(m)-\mathfrak{S}_\ell(u)\big)\right|^2 \\ &\ll \sum_{u\leq y}\left|\int_{\mathfrak{M}} J(\alpha,z)J_\ell(\alpha,z)e(u\alpha)\,\mathrm{d}\alpha - \sum_{z<m^\ell\leq 2^\ell z}\mathfrak{S}_\ell(u)\right|^2 \\ &\quad + \sum_{u\leq y}\left|\int_{\mathfrak{m}} J(\alpha,z)J_\ell(\alpha,z)e(u\alpha)\,\mathrm{d}\alpha\right|^2 := S_{\mathfrak{M}}(y,z)+S_{\mathfrak{m}}(y,z). \end{aligned} \tag{3.5}$$

Similarly, we have

$$F'_\ell(y,z) \ll S'_{\mathfrak{M}}(y,z) + S'_{\mathfrak{m}}(y,z), \tag{3.6}$$

where

$$S'_{\mathfrak{M}}(y,z) = \sum_{u\leq y}\left|\int_{\mathfrak{M}} J(\alpha,z)I_\ell(\alpha,z)e(u\alpha)\,\mathrm{d}\alpha - \sum_{z<m^\ell\leq 2^\ell z}\mathfrak{S}'_\ell(u)\right|^2$$

and

$$S'_{\mathfrak{m}}(y,z) = \sum_{u\leq y}\left|\int_{\mathfrak{m}} J(\alpha,z)I_\ell(\alpha,z)e(u\alpha)\,\mathrm{d}\alpha\right|^2.$$

We shall prove the following lemmas, from which, (3.1), (3.2), (3.5) and (3.6) the results of our two theorems follows.

Lemma 3.1. *For any $A>0$,there exists a $B_{\ell,1}(A)>0$ such that for $B\geq B_{\ell,1}(A)$, then*

$$S_{\mathfrak{m}}(y,z)\ll yz^{2/\ell}L^{-A}$$

holds for all $z^{1-1/(2\ell)}L^{B}\leq y\leq z$.

For any $A>0$,there exists a $B'_{\ell,1}(A)>0$ such that for $B\geq B'_{\ell,1}(A)$, then

$$S_{\mathfrak{m}}(y,z)\ll yz^{2/\ell}L^{-A}$$

holds for all $z^{1-1/\ell}L^{B}\leq y\leq z$.

Lemma 3.2. *For any $\delta>0$. Let $z^{\delta}<y\leq z$. Then we have*

$$S_{\mathfrak{M}}(y,z)\ll_{\delta,\ell,B} yz^{\frac{2}{\ell}}L^{-A} \text{ and } S'_{\mathfrak{M}}(y,z)\ll_{\delta,\ell,B} yz^{\frac{2}{\ell}}L^{-A}$$

holds for any $A>0$, where $B=(1+A)\max(6000\ell^3,2^{\ell}10^2\ell+c(\ell))$ with $c(\ell)$ is an absolute constant depends only on ℓ.

4. The minor arcs

In this section, we shall prove Lemma 3.1. Firstly, we have

$$S_{\mathfrak{m}}(y,z)\ll\int_{\mathfrak{m}}|J(\alpha,z)J_{\ell}(\alpha,z)|^2\,\mathrm{d}\alpha\ll z\sup_{\alpha\in\mathfrak{m}}|J_{\ell}(\alpha,z)|^2$$

by Bessel inequity. Then the classical results

$$J_{\ell}(\alpha,z)\ll z^{1/\ell}(\log z)^{-B}$$

holds for all $\alpha\in\mathfrak{m}$ and any$B\geq 0$ implies if $y\in(zL^{-B},z]$, one has

$$S_{\mathfrak{m}}(y,z)\ll z^{1+2/\ell}L^{-2B}\ll yz^{2/\ell}L^{-B}. \tag{4.1}$$

If $y\in(z^{1-1/(2\ell)}L^{B},zL^{-B}]$, then

$$\begin{aligned}S_{\mathfrak{m}}(y,z)&=\sum_{u\leq y}\left|\int_{\mathfrak{m}}J(\alpha,z)J_{\ell}(\alpha,z)e\left(u\alpha\right)\mathrm{d}\alpha\right|^2\\&=\int_{\mathfrak{m}}\mathrm{d}\beta J(\beta,z)J_{\ell}(\beta,z)\int_{\mathfrak{m}}J(-\alpha,z)J_{\ell}(-\alpha,z)\sum_{u\leq y}e\left(u(\alpha-\beta)\right)\mathrm{d}\alpha\\&\ll\int_{\mathfrak{m}}\mathrm{d}\beta\left|J(\beta,z)J_{\ell}(\beta,z)\right|\int_{\mathfrak{m}}\left|J(\alpha,z)J_{\ell}(\alpha,z)\right|\min\left(y,\frac{1}{\parallel\alpha-\beta\parallel}\right)\mathrm{d}\alpha.\end{aligned}$$

Splitting the unit interval in $H=\lfloor y\rfloor+1$ adjacent, disjoint intervals H_i of length H^{-1}, we obtain that

$$S_{\mathfrak{m}}(y,z)\ll\sum_{1\leq i,j\leq H}\sum\frac{y}{1+|i-j|}\iint_{\substack{\mathfrak{m}\cap H_i\\\mathfrak{m}\cap H_j}}\mathrm{d}\beta\mathrm{d}\alpha\left|J(\beta,z)J_{\ell}(\beta,z)J(\alpha,z)J_{\ell}(\alpha,z)\right|.$$

By cauchy inequity, we have

$$\begin{aligned}S_{\mathfrak{m}}(y,z)&\ll y\sum_{1\leq i\leq H}\left(\int_{\mathfrak{m}\cap H_i}\mathrm{d}\beta\left|J(\beta,z)J_{\ell}(\beta,z)\right|\right)^2\sum_{1\leq j\leq H}\frac{1}{1+|i-j|}\\&\ll y\log y\sum_{1\leq i\leq H}\left(\int_{\mathfrak{m}\cap H_i}\mathrm{d}\beta\left|J(\beta,z)\right|^2\int_{\mathfrak{m}\cap H_i}\mathrm{d}\beta\left|J_{\ell}(\beta,z)\right|^2\right)\\&\ll yL\int_{\mathfrak{m}}|J(\alpha,z)|^2\,\mathrm{d}\alpha\max_{1\leq i\leq H}\int_{\mathfrak{m}\cap H_i}\mathrm{d}\beta\left|J_{\ell}(\beta,z)\right|^2.\end{aligned}$$

For $1 \le i \le H$, there exist a $\beta = a/q + \lambda \in \mathfrak{m} \cap H_i$ with $(q,a) = 1$. Applying Gallaghers lemma (see [13, Lemma 1]) we have

$$\begin{aligned}
&\int_{\mathfrak{m}\cap H_i} \mathrm{d}\beta \, |J_\ell(\beta, z)|^2 \\
&\ll \int_{|\lambda| \le \frac{1}{H}} \mathrm{d}\lambda \left| \sum_{z < m^\ell \le 2^\ell z} \Lambda(m) e\left(\frac{a}{q} m^\ell\right) e\left(m^\ell \lambda\right) \right|^2 \\
&\ll \frac{1}{H^2} \int_{\mathbb{R}} \mathrm{d}x \left| \sum_{x \le m^\ell \le x + H/2} 1_{z < m^\ell \le 2^\ell z} \Lambda(m) e\left(\frac{a}{q} m^\ell\right) \right|^2 \\
&= \frac{1}{H^2} \int_{z - H/2}^{2^\ell z} \mathrm{d}x \left| \sum_{x \le m^\ell \le x + H/2} 1_{z < m^\ell \le 2^\ell z} \Lambda(m) e\left(\frac{a}{q} m^\ell\right) \right|^2 \\
&\ll \frac{1}{H^2} \left(\int_z^{2^\ell z - H/2} \mathrm{d}x \left| \sum_{x \le m^\ell \le x + H/2} \Lambda(m) e\left(\frac{a}{q} m^\ell\right) \right|^2 + H(H z^{\frac{1}{\ell} - 1})^2 \right) \\
&\ll \frac{1}{H^2} \sum_{j=0}^{\ell - 1} \int_{2^j z}^{2^{j+1} z} \mathrm{d}x \left| \sum_{x \le m^\ell \le x + H/2} \Lambda(m) e\left(\frac{a}{q} m^\ell\right) \right|^2 + H z^{\frac{2}{\ell} - 2}.
\end{aligned}$$

Then by Lemma 2.4 and notice that $y \le z L^{-B}$, one has

$$S_{\mathfrak{m}}(y,z) \ll y L (z^{\frac{2}{\ell} - 1} L^{-A-2} + z^{\frac{2}{\ell} - 1} L^{-B}) \int_0^1 |J(\alpha, z)|^2 \, \mathrm{d}\alpha \ll y z^{\frac{2}{\ell}} L^{-A}$$

holds for any $A > 0$ if $B \ge \max(B_{\mathfrak{m},\ell}(A), A) + 2$. Combining (4.1) and above, we get

$$S_{\mathfrak{m}}(y,z) \ll y z^{\frac{2}{\ell}} L^{-A}$$

holds for any $B \ge \max(B_{\mathfrak{m},\ell}(A), A) + 2$. Finally, using Lemma 2.5 substitute Lemma 2.4, it is not difficult to obtain the proof of the estimate of $S'_{\mathfrak{m}}(y,z)$.

5. The major arcs

In this section we consider the estimates $S_{\mathfrak{M}}(y,z)$ and $S'_{\mathfrak{M}}(y,z)$. For $S_{\mathfrak{M}}(y,z)$, notice that the definition of $B_\ell(q,a)$ (see Lemma 2.3), the fact

$$B_1(q,-a) = \sum_{h \in \mathbb{Z}_q^*} e\left(-\frac{ah}{q}\right) = \mu(q) \text{ if } (q,a) = 1,$$

(3.3) and (3.4) implies

$$\begin{aligned}
&\int_{\mathfrak{M}} J(\alpha,z)J_\ell(\alpha,z)e\left(u\alpha\right)\mathrm{d}\alpha-\mathfrak{S}_\ell(u)\sum_{z<m^\ell\le 2^\ell z}1\\
&=\sum_{q\le L^{B^2}}\sum_{a\in\mathbb{Z}_q^*}e\left(\frac{au}{q}\right)\int_{-\frac{L^{B^2}}{X}}^{\frac{L^{B^2}}{X}}\mathrm{d}\lambda J(\alpha,z)\left(J_\ell(\alpha,z)-\frac{B_\ell(q,a)}{\varphi(q)}I_\ell(\lambda,z)\right)e\left(u\lambda\right)\\
&+\sum_{q\le L^{B^2}}\sum_{a\in\mathbb{Z}_q^*}e\left(\frac{au}{q}\right)\frac{B_\ell(q,a)}{\varphi(q)}\int_{-\frac{L^{B^2}}{X}}^{\frac{L^{B^2}}{X}}\mathrm{d}\lambda I_\ell(\lambda,z)\left(J(\alpha,z)-\frac{\mu(q)}{\varphi(q)}I(\lambda,z)\right)e\left(u\lambda\right)\\
&+\left(\sum_{q\le L^{B^2}}\sum_{a\in\mathbb{Z}_q^*}e\left(\frac{au}{q}\right)\frac{\mu(q)B_\ell(q,a)}{\varphi(q)^2}\int_{-\frac{L^{B^2}}{X}}^{\frac{L^{B^2}}{X}}\mathrm{d}\lambda I(\lambda,z)I_\ell(\lambda,z)e\left(u\lambda\right)-\mathfrak{S}_\ell(u)\sqrt[\ell]{z}\right)\\
&=\int_{\mathfrak{M}}\mathrm{d}\alpha J(\alpha,z)R_\ell(\alpha,z)e\left(u\alpha\right)+\int_{\mathfrak{M}}\mathrm{d}\alpha R(\alpha)J_\ell(\alpha,z)e\left(u\alpha\right)\\
&+\left(\mathfrak{S}_\ell(u,L^{B^2})\int_{-\frac{L^{B^2}}{X}}^{\frac{L^{B^2}}{X}}\mathrm{d}\lambda I(\lambda,z)I_\ell(\lambda,z)e\left(u\lambda\right)-\mathfrak{S}_\ell(u)\sqrt[\ell]{z}\right):=R_1+R_2+R_3,
\end{aligned}$$

where

$$R(\alpha)=\sum_{m\le 2^\ell z}\Lambda(m)e\left(-m\alpha\right)-\frac{\mu(q)}{\varphi(q)}\sum_{m\le 2^\ell z}e\left(-m\lambda\right)$$

with $\alpha=a/q+\lambda$. Therefore, we have

$$S_{\mathfrak{M}}(y,z)=\sum_{u\le y}\left|R_1+R_2+R_3\right|^2\ll\sum_{u\le y}\left(\left|R_1\right|^2+\left|R_2\right|^2+\left|R_3\right|^2\right)$$

by Cauchy inequality. Similarly, we obtain that

$$S'_{\mathfrak{M}}(y,z)\ll\sum_{u\le y}\left(\left|R'_1\right|^2+\left|R'_2\right|^2+\left|R'_3\right|^2\right),$$

where

$$R'_1=\int_{\mathfrak{M}}\mathrm{d}\alpha J(\alpha,z)R'_\ell(\alpha,z)e\left(u\alpha\right)\ ,\ R'_2=\int_{\mathfrak{M}}\mathrm{d}\alpha R(\alpha)I_\ell(\alpha,z)e\left(u\alpha\right)$$

and

$$R'_3=\mathfrak{S}'_\ell(u,L^{B^2})\int_{-\frac{L^{B^2}}{X}}^{\frac{L^{B^2}}{X}}\mathrm{d}\lambda I(\lambda,z)I_\ell(\lambda,z)e\left(u\lambda\right)-\mathfrak{S}'_\ell(u)\sqrt[\ell]{z}.$$

We have firstly

$$\begin{aligned}
\sum_{u\le y}\left|R_1\right|^2&=\sum_{u\le y}\left|\sum_{q\le L^{B^2}}\sum_{a\in\mathbb{Z}_q^*}e\left(\frac{au}{q}\right)\int_{-\frac{L^{B^2}}{X}}^{\frac{L^{B^2}}{X}}\mathrm{d}\lambda J(\lambda,z)R_\ell(\alpha,z)e\left(u\lambda\right)\right|^2\\
&\le\sum_{u\le y}\left|\sum_{q\le L^{B^2}}\varphi(q)X\sup_{\alpha\in\mathfrak{M}}\left|R_\ell(\alpha,z)\right|X^{-1}L^{B^2}\right|^2\ll_B yz^{\frac{2}{\ell}}L^{-B}
\end{aligned}$$

by Lemma 2.1. Also

$$\begin{aligned}\sum_{u\le y}|R_2|^2&=\sum_{u\le y}\left|\sum_{q\le L^{B^2}}\sum_{a\in\mathbb{Z}_q^*}e\left(\frac{au}{q}\right)\int_{-\frac{L^{B^2}}{X}}^{\frac{L^{B^2}}{X}}\mathrm{d}\lambda J_\ell(\lambda,z)R(\alpha)e\left(u\lambda\right)\right|^2\\&\le\sum_{u\le y}\left|\sum_{q\le L^{B^2}}z^{\frac{1}{\ell}}\sum_{a\in\mathbb{Z}_q^*}\frac{|B(q,a)|}{\varphi(q)}\sup_{\alpha\in\mathfrak{M}}\left|\sum_{j=0}^{\lfloor(\log(2X))/\log 2\rfloor+1}R_1(-\alpha,2^j)\right|\frac{L^{B^2}}{X}\right|^2\\&\ll yz^{\frac{2}{\ell}}\left|\sum_{q\le L^{B^2}}L^{3B^2}\sup_{\alpha\in\mathfrak{M}}\max_{z\le 2X}|R_1(-\alpha,z)|\,X^{-1}\right|^2\ll yz^{\frac{2}{\ell}}L^{-B}.\end{aligned}$$

Note that $I(\lambda,z)\ll|\lambda|^{-1}$, Hua' inequity (see [14, Theorem 4])

$$\int_0^1|I_\ell(\lambda,z)|^{2^\ell}\,\mathrm{d}\lambda\ll_\ell z^{\frac{2^\ell-\ell}{\ell}}L^{c(\ell)},$$

where $c(\ell)$ is an absolute constant depend only on ℓ. Then the using of Hölder's inequality gives

$$\begin{aligned}&\int_{\frac{L^{B^2}}{X}<|\lambda|\le\frac12}\mathrm{d}\lambda\,|I(\lambda,z)I_\ell(\lambda,z)|\\&\ll\left(\int_{\frac{L^{B^2}}{X}<|\lambda|\le\frac12}\mathrm{d}\lambda\,|I(\lambda,z)|^{\frac{2^\ell}{2^\ell-1}}\right)^{\frac{2^\ell-1}{2^\ell}}\left(\int_0^1\mathrm{d}\lambda\,|I_\ell(\lambda,z)|^{2^\ell}\right)^{\frac{1}{2^\ell}}\\&\ll_\ell\left(L^{B^2}/X\right)^{(1-\frac{2^\ell}{2^\ell-1})\frac{2^\ell-1}{2^\ell}}\left(z^{\frac{2^\ell}{\ell}-1}L^{c(\ell)}\right)^{\frac{1}{2^\ell}}\ll z^{\frac{1}{\ell}}L^{-\frac{B-c(\ell)}{2^\ell}}.\end{aligned}$$

On the other hand,

$$\int_{-\frac12}^{\frac12}\mathrm{d}\lambda I(\lambda,z)I_\ell(\lambda,z)e\left(u\lambda\right)=z^{\frac{1}{\ell}}+O(1).$$

Setting $B\ge\max(2,c(\ell))$, we obtain that

$$\begin{aligned}\sum_{u\le y}|R_3|^2&=\sum_{u\le y}\left|\mathfrak{S}_\ell(u,L^B)\int_{|\lambda|\le\frac{L^B}{X}}\mathrm{d}\alpha I(\alpha,z)I_\ell(\alpha,z)e(u\alpha)-\mathfrak{S}_\ell(u)\sqrt[\ell]{z}\right|^2\\&\ll z^{\frac{2}{\ell}}\sum_{u\le y}\left|\mathfrak{S}_\ell(u,L^B)-\mathfrak{S}_\ell(u)\right|^2+z^{\frac{2}{\ell}}L^{-\frac{B-c(\ell)}{2^\ell}}\sum_{u\le y}|\mathfrak{S}_\ell(u)|^2.\end{aligned}$$

We can conclude from the above estimates that

$$S_{\mathfrak{M}}(y,z)\ll yz^{\frac{2}{\ell}}L^{-B}+z^{\frac{2}{\ell}}\sum_{u\le y}\left|\mathfrak{S}_\ell(u,L^B)-\mathfrak{S}_\ell(u)\right|^2+z^{\frac{2}{\ell}}L^{-\frac{B-c(\ell)}{2^\ell}}\sum_{u\le y}|\mathfrak{S}_\ell(u)|^2.$$

Similarly, we have

$$S'_{\mathfrak{M}}(y,z)\ll yz^{\frac{2}{\ell}}L^{-B}+z^{\frac{2}{\ell}}\sum_{u\le y}\left|\mathfrak{S}'_\ell(u,L^B)-\mathfrak{S}'_\ell(u)\right|^2+z^{\frac{2}{\ell}}L^{-\frac{B-c(\ell)}{2^\ell}}\sum_{u\le y}|\mathfrak{S}'_\ell(u)|^2.$$

Thus Lemma 3.2 is an immediate consequence of Lemma 6.1 and Lemma 6.3 of the following section by setting $B=(1+A)\max(6000\ell^3,2^\ell10^2\ell+c(\ell))$.

6. Singular series

In this section, we consider the estimate of the singular series. We firstly have the following lemma.

Lemma 6.1. *We have*

$$\sum_{u\le y}|\mathfrak{S}_\ell(u)|^2 \ll_\ell yL^{10\ell} \text{ and } \sum_{u\le y}|\mathfrak{S}'_\ell(u)|^2 \ll_\ell yL^{10\ell}.$$

Proof. Note that $0\le \varrho_\ell(p,u)\le \ell$ one has

$$\begin{aligned}\mathfrak{S}_\ell(u)&=\prod_{p|u}\frac{p-\varrho_\ell(p,u)}{p-1-\varrho_\ell(p,u)}\prod_p\left(1-\frac{\varrho_\ell(p,u)-1}{p-1}-\frac{\varrho_\ell(p,u)}{(p-1)^2}\right)\\&\ll_\ell\prod_{p|u,p>2\ell}\left(1+\frac{1}{p-1-\ell}\right)\prod_{p>2\ell}\left(1-\frac{\varrho_\ell(p,u)-1}{p-1}\right)\\&\ll_\ell\prod_{p|u,p>2\ell}\left(1+\frac{1}{p}\right)\prod_{p>2\ell}\left(1-\frac{\varrho_\ell(p,u)-1}{p}\right)\\&\ll_\ell\exp\left(\sum_{p|u,p>2\ell}\frac{1}{p}\right)\exp\left(\sum_{p>2\ell}\frac{1-\varrho_\ell(p,u)}{p}\right).\end{aligned}$$

Then by Lemma 2.3, we trivially have

$$\mathfrak{S}_\ell(u)\ll_\ell\exp\left(\log\log|u|+O(1)+4\ell\log\log(2|u|)+O_\ell(1)\right)\ll_\ell(\log(2|u|))^{5\ell}$$

holds for all $|u|\ge 1$. Similarly we have

$$\mathfrak{S}'_\ell(u)\ll_\ell(\log(2|u|))^{5\ell}$$

holds for all $|u|\ge 1$. Then the lemma is obvious. □

Lemma 6.2. *Let positive real number x and y sufficiently large. Then we have*

$$\sum_{u\le y}|\mathfrak{P}_\ell(u,x)-\mathfrak{S}_\ell(u,x)|^2\ll_\ell yx^{-1}\log^{\ell^2}x+x^{4\log x}$$

and

$$\sum_{u\le y}|\mathfrak{P}'_\ell(u,x)-\mathfrak{S}'_\ell(u,x)|^2\ll_\ell yx^{-1}\log^{\ell^2}x+x^{4\log x}.$$

Proof. Denote by

$$P(x)=\prod_{p\le x}p \text{ and } S(u,x)=\mathfrak{P}_\ell(u,x)-\mathfrak{S}_\ell(u,x).$$

Then, we have

$$S(u,x)=\sum_{\substack{q>x\\q|P(x)}}\frac{\mu(q)}{\varphi(q)}A(q,u)=\sum_{\substack{x<q\le V\\q|P(x)}}\frac{\mu(q)}{\varphi(q)}A(q,u)+\sum_{\substack{q>V\\q|P(x)}}\frac{\mu(q)}{\varphi(q)}A(q,u),$$

where $V > x$. Let $\lambda_x = \log^{-1} x > 0$. Then we have

$$\sum_{\substack{q>V\\ q|P(x)}} \frac{\mu(q)}{\varphi(q)} A(q,u) \ll \sum_{\substack{q>V\\ q|P(x)}} \frac{\mu(q)^2}{\varphi(q)} |A(q,u)| \le \frac{1}{V^{\lambda_x}} \sum_{q|P(x)} \frac{q^{\lambda_x}\mu(q)^2}{\varphi(q)} |A(q,u)|$$
$$= \frac{1}{V^{\lambda_x}} \prod_{p\le x} \left(1 + \frac{p^{\lambda_x}|A(p,u)|}{p-1}\right) \le \frac{1}{V^{\lambda_x}} \prod_{p\le x} \left(1 + \frac{|A(p,u)|}{p-1}\right).$$

Setting $V = \exp(\log^2 x)$, and notice that

$$A(p,u) = \begin{cases} \frac{p}{p-1}(\varrho_\ell(p,u)-1) - 1 & u \equiv 0 (\mathrm{mod} p) \\ \frac{p}{p-1}(\varrho_\ell(p,u)-1) + \frac{1}{p-1} & u \not\equiv 0 (\mathrm{mod} p) \end{cases} \le \frac{p\ell}{p-1}.$$

Then

$$\sum_{u\le y} \left| \sum_{q>V, q|P(x)} \frac{\mu(q)}{\varphi(q)} A(q,u) \right|^2 \ll y \frac{1}{x^2} \prod_{p\le x} \left(1 + \frac{\ell}{p}\right)^2 \ll y x^{-2} \log^{2\ell} x.$$

One the other hand,

$$\sum_{u\le y} \left| \sum_{\substack{x<q\le V\\ q|P(x)}} \frac{\mu(q)}{\varphi(q)} A(q,u) \right|^2 = \sum_{u\le y} \sum\sum_{\substack{x<q_1,q_2\le V\\ q_1,q_2|P(x)}} \frac{\mu(q_1)\mu(q_2)}{\varphi(q_1)\varphi(q_2)} A(q,u)\overline{A(q,u)}$$
$$= \sum_{u\le y} \sum_{\substack{x<q\le V\\ q|P(x)}} \frac{\mu(q)^2}{\varphi(q)^2} |A(q,u)|^2 + T_R(V,x),$$

where

$$T_R(V,x)$$
$$= \sum\sum_{\substack{x<q_1\ne q_2\le V\\ q_1,q_2|P(x)}} \frac{\mu(q_1)\mu(q_2)}{\varphi(q_1)^2\varphi(q_2)^2} \sum\sum_{\substack{a_i\in\mathbb{Z}_{q_i}^*\\ i=1,2}} \sum\sum_{\substack{h_j\in\mathbb{Z}_{q_j}^*\\ j=1,2}} e\left(\frac{a_1 h_1^\ell}{q_1} - \frac{a_2 h_2^\ell}{q_2}\right) \sum_{u\le y} e\left(\frac{a_1q_2 - a_2q_1}{q_1q_2}u\right)$$
$$\le \sum\sum_{\substack{x<q_1\ne q_2\le V\\ q_1,q_2|P(x)}} \frac{\mu(q_1)^2\mu(q_2)^2}{\varphi(q_1)^2\varphi(q_2)^2} \sum\sum_{\substack{a_i\in\mathbb{Z}_{q_i}^*\\ i=1,2}} \sum\sum_{\substack{h_j\in\mathbb{Z}_{q_j}^*\\ j=1,2}} \left(\sum_{\lfloor (q_1q_2)^{-1}y\rfloor q_1q_2 < u\le y} 1 \right)$$
$$\le \sum\sum_{\substack{x<q_1\ne q_2\le V\\ q_1,q_2|P(x)}} q_1q_2 \le \left(\sum_{x<q\le V, q|P(x)} q \right)^2 \ll V^4 \le x^{4\log x},$$

where the obviously fact $q_1q_2 \nmid (a_1q_2 - a_2q_1)$ has been used. On the other hand

$$\sum_{\substack{x<q\le V\\ q|P(x)}} \frac{\mu(q)^2}{\varphi(q)^2} |A(q,u)|^2 \ll x^{-1} \sum_{\substack{x<q\le V\\ q|P(x)}} \frac{\mu(q)^2 q}{\varphi(q)^2} |A(q,u)|^2 \ll x^{-1} \sum_{q|P(x)} \frac{\mu(q)^2 q}{\varphi(q)^2} |A(q,u)|^2$$
$$\ll x^{-1} \prod_{p\le x} \left(1 + \frac{p|A(p,u)|^2}{(p-1)^2}\right) \ll x^{-1} \log^{\ell^2} x.$$

Hence we obtain that

$$\sum_{u\le y}|S(u,x)|^2 \ll_\ell yx^{-1}\log^{\ell^2}x + x^{4\log x} + yx^{-2}\log^{2\ell}x \ll yx^{-1}\log^{\ell^2}x + x^{4\log x}.$$

Similarly, we obtain that

$$\sum_{u\le y}|\mathfrak{P}'_\ell(u,x)-\mathfrak{S}'_\ell(u,x)|^2 \ll_\ell yx^{-1}\log^{\ell^2}x + x^{4\log x}.$$

Which complete the proof of the lemma. □

Lemma 6.3. *Let $x\in[1,\exp(\sqrt[3]{\log y})]$. Then we have*

$$\sum_{u\le y}|\mathfrak{P}'_\ell(u,x)-\mathfrak{S}'_\ell(u)|^2 \ll_\ell yx^{-1/(2000\ell^2)}L^{2\ell}$$

and

$$\sum_{u\le y}|\mathfrak{P}_\ell(u,x)-\mathfrak{S}_\ell(u)|^2 \ll_\ell yx^{-1/(2000\ell^2)}L^{3\ell}.$$

Proof. By Lemma 6.1, Lemma 6.2 and Lemma 2.1 we have

$$\begin{aligned}\sum_{u\le y}|\mathfrak{P}'_\ell(u,x)-\mathfrak{S}'_\ell(u)|^2 &\ll \sum_{u\le\sqrt{y}}L^{10\ell}+\sum_{\sqrt{y}<u\le y}|\mathfrak{P}'_\ell(u,x)-\mathfrak{S}'_\ell(u)|^2\\ &\ll \sqrt{y}L^{10\ell}+yx^{-1}\log^{\ell^2}x+\sum_{\sqrt{y}<u\le y}|\mathfrak{S}'_\ell(u,x)-\mathfrak{S}'_\ell(u)|^2.\end{aligned}$$

Note that $x\in[1,\exp(\sqrt[3]{\log y})]$, we have

$$\begin{aligned}\sum_{u\le y}|\mathfrak{P}'_\ell(u,x)-\mathfrak{S}'_\ell(u)|^2 &\ll yx^{-1/2}+\sum_{j=0}^{\lfloor\frac{\log y}{\log 2}\rfloor+1}\sum_{2^j\sqrt{y}<u\le 2^{j+1}\sqrt{y}}|\mathfrak{S}'_\ell(u,x)-\mathfrak{S}'_\ell(u)|^2\\ &\ll_\ell yx^{-1/2}+yx^{-1/(2000\ell^2)}\log y \ll yx^{-1/(2000\ell^2)}L.\end{aligned}$$

On the other hand, we have

$$\mathfrak{P}_\ell(u,x)=\mathfrak{P}'_\ell(u,x)f_\ell(u,x),$$

where

$$f_\ell(u,x)=\prod_{p\mid u,p\le x}\left(1-\frac{1}{p-\varrho_\ell(p,u)}\right)^{-1}\prod_{p\le x}\left(1-\frac{\varrho_\ell(p,u)}{(p-1)(p-\varrho_\ell(p,u))}\right).$$

Let $x\to\infty$, then

$$\mathfrak{S}_\ell(u)=\mathfrak{S}'_\ell(u)f_\ell(u),$$

where

$$f_\ell(u)=\prod_{p\mid u}\left(1-\frac{1}{p-\varrho_\ell(p,u)}\right)^{-1}\prod_{p}\left(1-\frac{\varrho_\ell(p,u)}{(p-1)(p-\varrho_\ell(p,u))}\right).$$

It is easily seen that

$$f_\ell(u,x)\ll_\ell \log(|u|+2).$$

Hence

$$\begin{aligned}&\sum_{u\le y}|\mathfrak{P}_\ell(u,x)-\mathfrak{S}_\ell(u)|^2\\ &\quad\ll\sum_{u\le y}|\mathfrak{P}'_\ell(u,x)-\mathfrak{S}'_\ell(u)|^2 f_\ell(u,x)^2+\sum_{u\le y}|\mathfrak{S}'_\ell(u)|^2|f_\ell(u,x)-f_\ell(u)|^2.\end{aligned}$$

Therefore by Lemma 6.1 one has

$$\sum_{u\le y}|\mathfrak{P}_\ell(u,x)-\mathfrak{S}_\ell(u)|^2\ll L^2\sum_{u\le y}|\mathfrak{P}'_\ell(u,x)-\mathfrak{S}'_\ell(u)|^2+L^{2\ell}\sum_{u\le y}|1-f_\ell(u,x)/f_\ell(u)|^2.$$

On the other hand, if $x>\ell$, then

$$\begin{aligned}
&\sum_{u\le y}|f_\ell(u,x)/f_\ell(u)-1|^2\\
&\quad\ll\sum_{u\le y}\left|\prod_{p|u,p>x}\left(1-\frac{1}{p-\varrho_\ell(p,u)}\right)^{-1}\prod_{p>x}\left(1-\frac{\varrho_\ell(p,u)}{(p-1)(p-\varrho_\ell(p,u))}\right)-1\right|^2\\
&\quad=\sum_{u\le y}\left|\prod_{p|u,p>x}\left(1+\frac{1}{p}\right)\prod_{p>x}\left(1+O_\ell\left(\frac{1}{p^2}\right)\right)-1\right|^2\\
&\quad=\sum_{u\le y}\left|\exp\left(\sum_{p|u,p>x}\frac{1}{p}\right)\exp\left(O_\ell\left(\frac{1}{x\log x}\right)\right)-1\right|^2\\
&\quad=\sum_{u\le y}\left|\exp\left(1+O\left(\frac{\omega(u)}{x}\right)\right)\left(1+O_\ell\left(\frac{1}{x\log x}\right)\right)-1\right|^2,
\end{aligned}$$

where $\omega(u)$ is the number of distinct primes dividing u. Furthermore, it is well known that

$$0\le\omega(u)\ll\log(\log|u|+3).$$

Hence if $x>\max(\log(\log|u|+3),\ell)$, then

$$\sum_{u\le y}|f_\ell(u,x)/f_\ell(u)-1|^2\ll_\ell\frac{y}{x^2(\log x)^2}+\sum_{u\le y}\frac{\omega(u)^2}{x^2}\ll yx^{-1}.$$

Hence we obtain that

$$\sum_{u\le y}|\mathfrak{P}_\ell(u,x)-\mathfrak{S}_\ell(u)|^2\ll_\ell yx^{-1}L^{2\ell}+yx^{-1/(2000\ell^2)}L^3\ll yx^{-1/(2000\ell^2)}L^{3\ell}.$$

Thus we get the proof of the lemma. □

Acknowledgment

The author would like to thank Professor Zhi-Guo Liu for his consistent encouragement.

References

[1] Paul T. Bateman and Roger A. Horn. A heuristic asymptotic formula concerning the distribution of prime numbers. *Math. Comp.*, 16:363–367, 1962.

[2] A. F. Lavrik. On the distribution of k-twin primes. *Dokl. Akad. Nauk SSSR*, 132:1258–1260, 1960.

[3] Stephan Baier and Liangyi Zhao. Primes in quadratic progressions on average. *Math. Ann.*, 338(4):963–982, 2007.

[4] Timothy Foo and Liangyi Zhao. On primes represented by cubic polynomials. *Math. Z.*, 274(1-2):323–340, 2013.

[5] G. H. Hardy and J. E. Littlewood. Some problems of 'Partitio numerorum'; III: On the expression of a number as a sum of primes. *Acta Math.*, 44(1):1–70, 1923.

[6] Alessandro Zaccagnini. On the exceptional set for the sum of a prime and a kth power. *Mathematika*, 39(2):400–421, 1992.

[7] Koichi Kawada. On the asymptotic formula for the number of representations of numbers as the sum of a prime and a kth power. *Proc. Japan Acad. Ser. A Math. Sci.*, 69(8):283–286, 1993.

[8] Koichi Kawada. A zero density estimate for Dedekind zeta functions of pure extension fields. *Tsukuba J. Math.*, 22(2):357–369, 1998.
[9] Jörg Brüdern. Representations of natural numbers as the sum of a prime and a k-th power. *Tsukuba J. Math.*, 32(2):349–360, 2008.
[10] A. Perelli and A. Zaccagnini. On the sum of a prime and a kth power. *Izv. Ross. Akad. Nauk Ser. Mat.*, 59(1):185–200, 1995.
[11] C. Bauer. On the exceptional set for the sum of a prime and the k-th power of a prime. *Studia Sci. Math. Hungar.*, 35(3-4):291–330, 1999.
[12] Henryk Iwaniec and Emmanuel Kowalski. *Analytic number theory*, volume 53 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2004.
[13] P. X. Gallagher. A large sieve density estimate near $\sigma = 1$. *Invent. Math.*, 11:329–339, 1970.
[14] L. K. Hua. *Additive theory of prime numbers*. Translations of Mathematical Monographs, Vol. 13. American Mathematical Society, Providence, R.I., 1965.

Department of Mathematics, East China Normal University, 500 Dongchuan Road, Shanghai 200241, PR China
E-mail address: nianhongzhou@outlook.com